S know the facts

prescription pain relievers (opioids)? What are the risks of misusing

Talk with your health care provider about all the prescription drugs you take.

- Mental confusion
- 000 Drowsiness
- Changes in brain that build tolerance, leading to dependence

linfectious diseases such as HIV, Hepatitis B and Hepatitis C if injected

WITHDRAWAL SYMPTOMS:

Restlessness, muscle and bone pain, insomnia, diarrhea, vomiting, cold flashes with goose bumps, kicking movements, severe craving that can lead to relapse

Euphoria, warm flushing, poor nutrition, fatal overdose TOTAL BODY EFFECTS:

PREGNANCY AND INFANTS: Miscarriage, low birth weight, developmental delays, addiction in infants

0 000 0 0 Cramping Nausea HIV and other sexually transmitted infections from unsafe sex while intoxicated Slow, irregular breathing that can be fatal Increased and/or decreased blood pressure Constipation

Symptoms of overdose

WEAK PULSE DROWSINESS LOW BLOOD PRESSURE COMA BLUE NAILS AND LIPS MUSCLE SPASMS DISORIENTATION VOMITING NON RESPONSIVE SEIZURES SMALL PUPILS NO BREATHING OR OLD/CLAMMY SHALLOW BREATHING SKIN DRY MOUTH

Need help?

For free referrals and emotional support:

Teen Link

1-866-833-6546

Washington Recovery Help Line 1-866-789-1511

Quick facts

- 75 percent of ALL prescription drug overdoses drug overauses are from opioids.
 - O Prescription opioids are pain. medications that relieve
 - 0 They can be addictive if not used as directed by your doctor.
 - 0 More people die from opioid overdoses than from any other drug.



Nearly 17,000 people die each year from prescription pain reliever overdoses.

Lock up your meds and safely discard them. Visit: www.TakeBackYourMeds.org



